

## On Generalizations of Extending Modules

FATİH KARABACAK

*Anadolu University, Education Faculty, Department of Mathematics, 26470, Eskisehir, Turkey*

*e-mail: fatihkarabacak@anadolu.edu.tr*

**ABSTRACT.** A module  $M$  is said to be SIP-extending if the intersection of every pair of direct summands is essential in a direct summand of  $M$ . SIP-extending modules are a proper generalization of both SIP-modules and extending modules. Every direct summand of an SIP-module is an SIP-module just as a direct summand of an extending module is extending. While it is known that a direct sum of SIP-extending modules is not necessarily SIP-extending, the question about direct summands of an SIP-extending module to be SIP-extending remains open. In this study, we show that a direct summand of an SIP-extending module inherits this property under some conditions. Some related results are included about  $C_{11}$  and SIP-modules.

### 1. Introduction

Throughout this paper all rings are associative with unity and  $R$  always denotes such a ring. Modules are unital and for an abelian group  $M$ , we use  $M_R$  (resp.  ${}_R M$ ) to denote a right (resp. left)  $R$ -module. Let  $M$  be a  $R$ -module and  $N$  a submodule of  $M$ . We use  $N \leq_e M$  and  $N \leq_d M$  to denote that  $N$  is essential in  $M$  and  $N$  is a direct summand of  $M$ , respectively. Moreover we use  $End(M_R)$  and  $r(m)$  to denote the ring of endomorphisms of  $M$  and the right annihilator in  $R$  of an element  $m$  in  $M$ , i.e.,  $r(m) = \{r \in R : m.r = 0\}$ . Recall that a ring is called *Abelian* if every idempotent is central. For any unexplained terminology please see [1] and [5].

A module  $M_R$  has the *Summand Intersection Property*, SIP, if the intersection of every pair of direct summands of  $M_R$  is a direct summand of  $M_R$ . The study of modules having SIP was motivated by the following result of Kaplansky [7]: every free module over any principal ideal domain has SIP. The Summand Intersection Property has been studied by many authors (see e.g. [2], [3], [6], [8] and [17].)

Recall that a module  $M$  is called an *extending module* (or a *CS-module*) if every submodule is essential in a direct summand of  $M$ . In [5] and [11], extending modules were studied in detail.

The concept of  $C_{11}$ -modules was introduced in [15] as a generalization of extending modules. A module  $M$  is called  $C_{11}$ -module (or satisfies  $C_{11}$ )[15] if every

---

Received August 3, 2008; accepted November 4, 2008.

2000 Mathematics Subject Classification: 16D10, 16D15.

Key words and phrases: SIP-extending modules, summand intersection property, extending modules.

submodule of  $M$  has a complement which is a direct summand. It is known that a direct summand of a  $C_{11}$ -module is not a  $C_{11}$ -module, in general (see [16, Exercise 4]). A module is called a  $C_{11}^+$ -module if its every direct summand is a  $C_{11}$ -module [15]. In this paper we further the study of SIP-extending modules and we show that if  $M$  is a  $C_{11}$ -module which is also SIP-extending then every direct summand of  $M$  is a  $C_{11}$ -module, i.e.,  $M$  is  $C_{11}^+$ -module (see Proposition 7). In the main result we show that if  $M$  is an SIP-extending module such that  $End(M_R)$  is Abelian then every direct summand of  $M$  is SIP-extending.

## 2. SIP-extending modules

In [9], a module  $M$  is called an *SIP-extending module* provided that the intersection of every pair of direct summands of  $M$  is essential in a direct summand of  $M$ . We say a ring  $R$  is a right *SIP-extending ring* if the module  $R_R$  is an SIP-extending module, i.e., for every pair of idempotents  $e, c$  in  $R$  there exists  $g^2 = g \in R$  such that  $eR \cap cR$  is essential in  $gR$ . Examples of SIP-extending modules include every extending (hence every injective) module, every uniform module, every semisimple module and every module having the SIP (e.g. any Baer module [13]). The concept of an SIP-extending module is a proper generalization of both SIP-modules and extending modules, as shown by the following examples.

**Example 1.** Let  $F$  be any field and  $V$  be a  $F$ -vector space with  $dimV_F \geq 2$ . Let

$$R = \left\{ \begin{bmatrix} a & v \\ 0 & a \end{bmatrix} : a \in F, v \in V \right\},$$

be the trivial extension of  $F$  by  $V$ . Then  $R$  is a right SIP-extending ring however since  $dimV_F \geq 2$ ,  $R$  is not right extending ring.

**Example 2**([4, Exercise 1.5]). Let  $F$  be a field and

$$T = \left\{ \begin{bmatrix} a & x & 0 & 0 \\ 0 & b & 0 & 0 \\ 0 & 0 & b & y \\ 0 & 0 & 0 & a \end{bmatrix} : a, b, x, y \in F \right\}.$$

Let

$$e = e^2 = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

and

$$c = c^2 = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Then  $eT \cap cT$  is nilpotent. Hence  $eT \cap cT$  is not a direct summand of  $T$ . It follows that  $T$  does not have SIP. However it is a right SIP-extending ring.

It is well known that every direct summand of SIP-modules is an SIP-module and every direct summand of extending modules is an extending module. This result led us to the following question.

Question: Let  $M$  be an SIP-extending module and  $N$  be a direct summand of  $M$ . Is  $N$  an SIP-extending module?

In [9] we have provided a positive answer to the direct summand question under the condition that the summand is the unique closure of each of its essential submodules.

**Proposition 3**([9, Lemma 6]). *Let  $M$  be an SIP-extending module, and let  $N$  be a direct summand of  $M$ . Suppose that  $N$  is the unique closure in  $M$  of any of its essential submodules. Then  $N$  is also an SIP-extending module.*

**Definition 4**([14]). Let  $M$  be a module. If every submodule has a unique closure in  $M$  then  $M$  is called UC-module.

**Proposition 5.** *Let  $M$  be a UC-module. Then  $M$  has SIP if and only if  $M$  is SIP-extending.*

*Proof.* It is clear that if  $M$  has SIP then  $M$  is SIP-extending. Conversely, let  $S_1$  and  $S_2$  be direct summands of  $M$ . Then by hypothesis  $S_1 \cap S_2 \leq_e P$  for some  $P \leq_d M$ . By the main theorem in [14], intersection of two closed submodules is closed hence  $S_1 \cap S_2 = P$ . Thus  $M$  has SIP.  $\square$

The following lemma is proved in [9, Theorem 8].

**Lemma 6.** *Let  $M$  be a  $C_{11}$ -module and  $E$  be a submodule of  $M$ . If for every direct summand  $D$  of  $M$ ,  $E \cap D$  is essential in a direct summand of  $E$  then  $E$  is a  $C_{11}$ -module.*

**Lemma 7.** *Let  $M$  be a  $C_{11}$ -module. If  $M$  is SIP-extending then every direct summand of  $M$  is  $C_{11}$  (i.e.,  $M$  has  $C_{11}^+$ ).*

*Proof.* By Lemma 6 and the definition of SIP-extending.  $\square$

Recall that  $R$  is said to Abelian if every idempotent of  $R$  is central. Note that every finite dimension module has an Abelian endomorphism ring by [11]. We have the following result for SIP-extending Abelian rings.

**Proposition 8.** *Let  $R$  be an Abelian ring then*

- i)  $R$  is SIP-extending (SIP) if and only if  $R[x]$  is SIP-extending (SIP).
- ii)  $R$  is SIP-extending (SIP) if and only if  $R[[x]]$  is SIP-extending (SIP).

*Proof.* Since  $R$  is an Abelian ring, the result follows by [10, Lemma 8].  $\square$

Next, we provide an answer to the direct summand question for an SIP-extending module under another special condition. The result shows that a fully

invariant direct summand of an SIP-extending module inherits the property. It also completes the sufficiency part of [9, Theorem 11] in which only the necessity was established.

**Theorem 9.** *Let  $M = \bigoplus_{i \in I} M_i$  be a direct sum of fully invariant submodules  $M_i$  of  $M$  where  $I$  is an index set. Then  $M$  is an SIP-extending module if and only if  $M_i, \forall i \in I$  is an SIP-extending module.*

*Proof.* Let  $M$  be an SIP-extending module and  $M_i$  be a fully invariant direct summand of  $M$ . If  $L$  and  $K$  are direct summand of  $M_i$  then there exist  $P \leq_d M$  ( $M = P \oplus Q$ , for some  $Q \leq M$ ) such that  $L \cap K \leq_e P$ . Since  $M_i$  is a fully invariant direct summand of  $M$  and  $M = P \oplus Q$  then  $M_i = (M_i \cap P) \oplus (M_i \cap Q)$ . Therefore  $L \cap K \leq_e M_i \cap P \leq_d M_i$ . So  $M_i$  is an SIP-extending module. Converse follows from [9, Theorem 11]. We include a brief outline for the convenience of the reader. Let  $S$  be any direct summand of  $M$ . So  $S = \bigoplus (S \cap M_i)$ . Now let  $S, T$  be direct summands of  $M$ . Hence,  $S \cap T = \bigoplus [(S \cap M_i) \cap (T \cap M_i)]$ . Therefore, there exists a direct summand  $K_i$  of  $M_i$  which contains  $(S \cap M_i) \cap (T \cap M_i)$  as an essential submodule.  $\square$

**Corollary 10.**  *$M_R$  is an SIP-extending module such that  $End(M_R)$  is Abelian. Then every direct summand of  $M$  is SIP-extending.*

*Proof.* Let  $M$  be an SIP-extending module and  $M_1$  be a direct summand of  $M$ . Since  $End(M_R)$  is Abelian  $M_1$  is a fully invariant submodule of  $M$ . By Theorem 9,  $M_1$  is an SIP-extending module.  $\square$

**Definition 11.** Let  $M$  be a  $R$ -module.  $M$  is said to be multiplication module if for each  $X \leq M$  there exists  $A_R \leq R_R$  such that  $X = MA$

**Corollary 12.** *Let  $M$  be an SIP-extending module, then any direct summand of  $M$  is SIP-extending if  $M$  satisfies any of the following conditions.*

- (i)  $M_R = R_R$  and  $R$  is Abelian.
- (ii)  $M$  is a multiplication module and  $R$  is commutative.

*Proof.* (i) Immediate by Corollary 10.

(ii) Assume that  $M$  is multiplication module and  $R$  is commutative. Note that every submodule of a multiplication module is fully invariant. Now Theorem 9 yields the result.  $\square$

Recall that a module  $M$  satisfies the  $C_3$  condition whenever  $K, L$  are direct summand of  $M$  with  $K \cap L = 0$  then  $K \oplus L \leq_d M$  (see [11]). Note that the  $\mathbb{Z}$ -module ( $\mathbb{Z} \oplus \mathbb{Z}$ ) is an SIP-extending which does not satisfy the  $C_3$  condition (see, for example [3]). Now we provide an example which shows that a module satisfying the  $C_3$  property does not have to be SIP-extending either.

**Example 13.** Let  $F$  be a field and  $R = \begin{pmatrix} F & F \\ 0 & F \end{pmatrix}$  be the ring of upper triangular matrices over  $F$ ,  $N = \begin{pmatrix} 0 & F \\ 0 & F \end{pmatrix}$  and  $L = \begin{pmatrix} F & F \\ 0 & 0 \end{pmatrix}$  left ideals of  $R$  and  $M =$

$R/L$ . Let  $U = N \oplus M$ . Then  ${}_R U$  satisfies the  $C_3$  condition and is a UC-module but does not have the SIP as a left  $R$ -module. By Proposition 5,  $U$  is not SIP-extending as a left  $R$ -module.

We conclude this paper with some results for  $C_{11}$ -modules. Recall that a module  $M$  is said to satisfy the full (finite) exchange property if for any module  $G$  and any two direct sum decompositions  $G = M' \oplus N = \bigoplus_{i \in I} A_i$  where  $M' \cong M$  and  $I$  is any (finite) index set, there are submodules  $B_i$  of  $A_i$ ,  $i \in I$ , such that  $G = M' \oplus (\bigoplus_{i \in I} B_i)$ .

It was shown in [12] that every quasi-continuous module (i.e., an extending module with  $C_3$  condition) satisfies the full exchange property whenever it satisfies the finite exchange property. We can weaken the extending condition ( $C_1$ ) to  $C_{11}$  under an additional chain condition.

**Theorem 14.** *Let  $M_R$  be a  $C_{11}$ -module which satisfies  $C_3$  condition. If  $M$  has ACC on  $r(m)$ ,  $m \in M$  then the finite exchange property of  $M$  implies the full exchange property.*

*Proof.* By [15, Lemma 4.6 (a)] and [11, Theorem 2.17]  $M$  has a decomposition into indecomposable submodules. This yields the full exchange property.  $\square$

**Corollary 15.** *Let  $R$  be a right Noetherian ring and  $M_R$  be a  $C_{11}$ -module which has  $C_3$  then the finite exchange property implies the full exchange property for  $M$ .*

*Proof.* It follows from Theorem 14.  $\square$

## References

- [1] F. W. Anderson and K. R. Fuller, *Rings and Categories of Modules*, Springer-Verlag, 1974.
- [2] D. M. Arnold and J. Hausen, *A characterization of modules with the summand intersection property*, *Comm. Algebra*, **18**(1990), 519-528.
- [3] G. F. Birkenmeier, F. Karabacak and A. Tercan, *When is the SIP (SSP) property inherited by free modules*, *Acta Math. Hungar.*, **112**(2006), 103-106.
- [4] G. F. Birkenmeier, J. Y. Kim and J. K. Park, *When is the CS condition hereditary?*, *Comm. Algebra*, **27**(1999), 3785-3885.
- [5] N. V. Dung, D. V. Huynh, P. F. Smith and R. Wisbauer, *Extending Modules*, Longman, 1990.
- [6] J. Hausen, *Modules with the summand intersection property*, *Comm. Algebra*, **17**(1989), 135-148.
- [7] I. Kaplansky, *Infinite Abelian Groups*, University of Michigan Press, 1969.
- [8] F. Karabacak and A. Tercan, *Matrix rings with the summand intersection property*, *Czech. Math. J.*, **53**(2003), 621-626.
- [9] F. Karabacak and A. Tercan, *On modules and matrix rings with SIP-extending*, *Taiwanese J. Math.*, **11**(2007), 1037-1044.

- [10] N. K. Kim and Y. Lee, *Armendariz rings and reduced rings*, Journal of Algebra, **223**(2000), 477-488.
- [11] S. H. Mohamed and B. J. Müller, *Continuous and Discrete Modules*, Cambridge University Press, 1990.
- [12] K. Oshiro and S. T. Rizvi, *Exchange property of quasi-continuous modules with the finite exchange property*, Osaka J. Math., **33**(1996), 217-234.
- [13] S. T. Rizvi and C. S. Roman, *Baer and quasi-baer modules*, Comm. Algebra, **32**(2004), 103-123.
- [14] P. F. Smith, *Modules for which every submodule has a unique closure*, Ring Theory (S.Jain and S.T. Rizvi eds.) New Jersey, World Scientific, (1992), 302-317.
- [15] P. F. Smith and A. Tercan, *Generalizations of CS-modules*, Comm. Algebra, **21**(1993), 1809-1847.
- [16] P. F. Smith and A. Tercan, *Direct summands of modules which satisfy  $(C_{11})$* , Algebra Colloq., **11**(2004), 231-237.
- [17] G. V. Wilson, *Modules with the summand intersection property*, Comm. Algebra, **14**(1986), 21-38.