

Lichen Records from Sarıçık Mountain in Southern Giresun Province, Turkey

Demet CANSARAN DUMAN*, Ender YURDAKULOL

University of Ankara, Faculty of Science, Department of Biology, Botany Section, 06100 Tandoğan, Ankara - TURKEY

Received: 16.11.2005

Accepted: 28.02.2007

Abstract: Herein, we report 87 lichen species from 11 localities in the Sarıçık Mountains, Giresun, Turkey, of which 48 are new records for Giresun Province.

Key Words: Lichens, flora, Giresun, Turkey

Giresun İlinin Güneyinde bulunan Sarıçık Dağından Liken Kayıtları, Türkiye

Özet: Sarıçık Dağı'nda (Giresun) 11 lokaliteden 87 lichen türü kaydedilmiştir. Bunlardan 48'i Giresun ili için yeni kayıtdır.

Anahtar Sözcükler: Likenler, flora, Giresun, Türkiye

Introduction

Although several detailed studies on the lichen flora of Turkish provinces have recently been published (Güvenç, 2002; Yıldız et al., 2002; Yıldız & John, 2002; Çobanoğlu & Akdemir, 2004; Güvenç & Öztürk, 2004; Öztürk et al., 2005), knowledge of the lichen flora in some regions is still incomplete. Currently, approximately 152 species have been published from Giresun Province (Steiner, 1909; Küçük, 1990; Özgen et al., 2002; Süleyman et al., 2002; John & Breuss, 2004; Kinalioğlu, 2005; Yazıcı & Aslan, 2005). The present study was conducted in a semi-natural area and contributes to our knowledge of the chorology of lichens in Turkey. As environmental and ecological indicators, lichens helped to characterise the habitats that exist within the study area.

The study area is situated at lat 40°12'-40°16'N, long 38°39'-38°53'E. According to the grid system in *Flora of Turkey* the study area is in the A4 square (Davis, 1965). Sarıçık Mountain has a mean altitude of 2411 m and is bordered by Şebinkarahisar and Giresun to the north-

west, Alucra to the north, Şiran to the east, Refahiye to the south, and Zara to the south-west. The localities are shown in Figure 1.

The study area is located in the Xero-Euxine sub-province of the Euro-Siberian region in Turkey. In collection sites 2 and 5 of the study area, steppic vegetation was found, especially on uncultivated land and slopes. Within the study area the vegetation is under the influence of heavy anthropogenic factors and the effect of these factors could be observed in different plant associations of the Irano-Turanian phytogeographic region, such as at collection site 7.

In the study area, deciduous forest vegetation prevails mainly at lower altitudes and forms a belt surrounding the pine forests. The floristic composition here includes *Quercus petraea* (Mattuschka) Liebl. subsp. *iberica* (Steven ex Bieb.) Krassiln., *Q. pubescens* Willd., *Q. infectoria* Oliver., and *Carpinus betulus* L. populations. This vegetation type is represented by collection sites 1, 3, 10, and 11.

* E-mail: cansaran@science.ankara.edu.tr




Figure 1. The study area.

The pine forests have 2 dominant species, *Pinus sylvestris* L. and *Abies nordmanniana* (Stev.) Spach. subsp. *nordmanniana*. These pine forests are located at the higher altitudes of the study area and so reduced impact of the destruction caused by settlements can be seen in the floristic structure. Pine forests were studied at collection sites 4, 6, and 8.

The other companion species of the floristic structure on the tree level are *Corylus avellana* L., *Salix alba* L., and *Populus tremula* L. This vegetation type is represented by collection site 9.

According to meteorological data, the study area is under the influence of a semi-arid, very cold subtype of the Mediterranean climate (Akman, 1990).

Materials and Methods

The lichen specimens with their substrates were collected between 2001 and 2002 from 11 localities. The collected specimens were identified using flora books (Purvis et al., 1992; Wirth, 1995). The specimens were prepared as herbarium material and stored in the ANK herbarium.

Results

In all, 87 lichen taxa from 19 families were identified from 11 localities in the Sarıçek Mountains (Şebinkarahisar, Giresun), of which 48 taxa are new records for the province of southern Giresun. The taxa are listed in alphabetical order, followed by localities and substrates. Author names are given according to Brummitt & Powell (1992). New species for Giresun province are indicated by an asterisk (*).

List of Taxa

**Acarospora cervina* A.Massal.

Giresun: Şebinkarahisar; Körülü Valley, lat 40°14'N, long 38°39'E, alt. 1750 m, on calcareous rocks, 12.09.2001, ANK 48.

**Anaptychia ciliaris* (L.) Körb. ex A.Massal.

Giresun: Şebinkarahisar; Kirtepe-Yaygış, lat 40°13'N, long 38°40'E, alt. 2150 m, on siliceous rocks, 11.09.2001, ANK 49.

**Anaptychia setifera* Mereschk. ex Räsänen

Giresun: Şebinkarahisar; Kutluca Plateau, lat 40°11'N, long 38°48'E, alt. 2000 m, on *Abies nordmanniana* subsp. *nordmanniana*, 11.09.2001, ANK 50.

**Aspicilia calcarea* (L.) Mudd

Giresun: Şebinkarahisar; Körülü Valley, lat 40°14'N, long 38°39'E, alt. 1750 m, on calcareous rocks, 12.09.2001, ANK 51.

Aspicilia cinerea (L.) Körb.

Giresun: Şebinkarahisar; around Çakılıkaya, lat 40°12'N, long 38°50'E, alt. 2050 m, on calcareous rocks, 17.07.2002; Kirtepe-Yaygış, lat 40°13'N, long 38°40'E, alt. 2150 m, on calcareous rocks, 11.09.2001, ANK 52.

**Aspicilia contorta* (Hoffm.) Kremp.

Giresun: Şebinkarahisar; around Çakılıkaya, lat 40°12'N, long 38°50'E, alt. 2050 m, on siliceous rocks, 17.07.2002, ANK 53.

**Bellemera cinereorufescens* (Ach.) Clauzade & Cl.Roux

Giresun: Şebinkarahisar; Körülü Valley, lat 40°14'N, long 38°39'E, alt. 1750 m, on calcareous rocks, 12.09.2001, ANK 54.

**Bryoria chalybeiformis* (L.) Brodo & D.Hawksw.

Giresun: Şebinkarahisar; Slope 4 km south of İncedere Plateau, lat 40°11'N, long 38°53'E, alt. 1750 m, on *Abies nordmanniana* subsp. *nordmanniana*, 11.09.2001, ANK 03.

**Bryoria fuscescens* (Gyeln.) Brodo & D.Hawksw.

Giresun: Şebinkarahisar; Kabaktepe, lat 40°13'N, long 38°38'E, alt. 1950 m, on *Pinus syvestris*, 11.09.2001, ANK 55.

**Bryoria nadvornikiana* (Gyeln.) Brodo & D.Hawksw.

Giresun: Şebinkarahisar; Kutluca Plateau, lat 40°11'N, long 38°48'E, alt. 2000 m, on *Abies nordmanniana* subsp. *nordmanniana*, 11.09.2001, ANK 04.

**Caloplaca biatorina* (A.Massal.) J.Steiner

Giresun: Şebinkarahisar; around Çakılıkaya, lat 40°12'N, long 38°50'E, alt. 2050 m, on calcareous rocks, 17.07.2002, ANK 56.

**Caloplaca cerina* (Ehrh. ex Hedw.) Th.Fr.

Giresun: Şebinkarahisar; Yıldızlarla Hill, lat 40°14'N, long 38°48'E, alt. 1750 m, on *Populus tremula*, 11.09.2001; Kutluca Plateau, lat 40°11'N, long 38°48'E, alt. 2000 m, on mosses, 11.09.2001, ANK 57.

Caloplaca flavescens (Huds.) J.R.Laundon

Giresun: Şebinkarahisar; Kirtepe-Yaygış, lat 40°13'N, long 38°40'E, alt. 2150 m, on calcareous rocks, 11.09.2001, ANK 58.

Caloplaca holocarpa (Hoffm. ex Ach.) A.E.Wade

Giresun: Şebinkarahisar; on top of and around Büyükkal Hill, lat 40°19'N, long 38°40'E, alt. 2000 m, on *Abies nordmanniana* subsp. *nordmanniana*, 12.09.2001, ANK 46.

**Caloplaca luteoalba* (Turner) Th.Fr.

Giresun: Şebinkarahisar; on top of and around Büyükkal Hill, lat 40°19'N, long 38°40'E, alt. 2000 m, on *Abies nordmanniana* subsp. *nordmanniana*, 12.09.2001, ANK 47.

**Caloplaca variabilis* (Pers.) Müll.Arg.

Giresun: Şebinkarahisar; Kutluca Plateau, lat 40°11'N, long 38°48'E, alt. 2000 m, on limestone rocks, 11.09.2001, ANK 59.

Candelariella vitellina (Hoffm.) Müll.Arg.

Giresun: Şebinkarahisar; Körülü Valley, lat 40°14'N, long 38°39'E, alt. 1750 m, on siliceous rocks, 12.09.2001, ANK 60.

**Candelariella xanthostigma* (Pers. ex Ach.) Lettau

Giresun: Şebinkarahisar; on top of and around Büyükkal Hill, lat 40°19'N, long 38°40'E, alt. 2000 m, on *Populus tremula*, 12.09.2001, ANK 05.

**Carbonea vitellinaria* (Nyl.) Hertel

Giresun: Şebinkarahisar; Kutluca Plateau, lat 40°11'N, long 38°48'E, alt. 2000 m, on *Candelariella vitellina*, 11.09.2001, ANK 61.

Cetraria islandica (L.) Ach.

Giresun: Şebinkarahisar; Kirtepe-Yaygış, lat 40°13'N, long 38°40'E, alt. 2150 m, on soil, 11.9.2001, ANK 13.

**Cetrelia cetrariooides* (Duby) W.L.Culb. & C.F.Culb.

Giresun: Şebinkarahisar; slope 4 km south of İncedere Plateau, lat 40°11'N, long 38°53'E, alt. 1750 m, on *Salix alba*, 11.09.2001, ANK 62.

**Chaenotheca chrysoccephala* (Ach.) Th.Fr.

Giresun: Şebinkarahisar; around Çakıkaya, lat 40°12'N, long 38°50'E, alt. 2050 m, on *Abies nordmanniana* subsp. *nordmanniana*, 17.07.2002, ANK 02.

**Chrysothrix candelaris* (L.) J.R.Laundon

Giresun: Şebinkarahisar; on top of and around Büyükçal Hill, lat 40°19'N, long 38°40'E, alt. 2000 m, on *Abies nordmanniana* subsp. *nordmanniana*, 12.09.2001, ANK 01.

**Cladonia ciliata* Stirt.

Giresun: Şebinkarahisar; Yıldızlarla Hill, lat 40°14'N, long 38°48'E, alt. 1750 m, on the mosses, 11.09.2001, ANK 63.

Cladonia coniocraea (Flörke) Spreng.

Giresun: Şebinkarahisar; Menekşeli-Ovacık, lat 40°16'N, long 38°46'E, alt. 1900 m, on soil, 11.09.2001, ANK 64.

Cladonia fimbriata (L.) Fr.

Giresun: Şebinkarahisar; on top of and around Büyükçal Hill, lat 40°19'N, long 38°40'E, alt. 2000 m, on soil, 12.09.2001, ANK 06.

Cladonia pyxidata (L.) Hoffm.

Giresun: Şebinkarahisar; around Çakıkaya, lat 40°12'N, long 38°50'E, alt. 1950 m, on soil, 11.09.2001, ANK 65.

Dermatocarpon intestiniforme (Körb.) Hasse

Giresun: Şebinkarahisar; Yıldızlarla Hill, lat 40°14'N, long 38°48'E, alt. 1750 m, on calcareous rocks, 11.09.2001, ANK 66.

Dermatocarpon luridum (With.) J.R.Laundon

Giresun: Şebinkarahisar; Çakıkaya, lat 40°12'N, long 38°50'E, alt. 2050 m, on calcareous rocks, 11.09.2001, ANK 67.

Dermatocarpon miniatum (L.) Mann

Giresun: Şebinkarahisar; around Çakıkaya, lat 40°12'N, long 38°50'E, alt. 2050 m, on calcareous rocks, 11.09.2001, ANK 68.

**Dimelaena oreina* (Ach.) Norman

Giresun: Şebinkarahisar; Köroğlu Valley, lat 40°14'N, long 38°39'E, alt. 1750 m, on siliceous rocks, 12.09.2001, ANK 69.

**Evernia divaricata* (L.) Ach.

Giresun: Şebinkarahisar; Kabaktepe Hill, lat 40°13'N, long 38°38'E, alt. 1950 m, on the *Abies nordmanniana* subsp. *nordmanniana*, 11.09.2001; Köroğlu Valley, lat 40°14'N, long 38°39'E, alt. 1750 m, on siliceous rocks, 12.09.2001, ANK 15.

**Evernia prunastri* (L.) Ach.

Giresun: Şebinkarahisar; Yıldızlarla Hill, lat 40°14'N, long 38°48'E, alt. 1750 m, on *Pinus sylvestris*, 11.09.2001, ANK 70.

Flavoparmelia caperata (L.) Hale

Giresun: Şebinkarahisar; Kabaktepe Hill, lat 40°13'N, long 38°38'E, alt. 1950 m, on *Abies nordmanniana* subsp. *nordmanniana*, 11.09.2001, ANK 71.

Graphis scripta (L.) Ach.

Giresun: Şebinkarahisar; Menekşeli-Ovacık, lat 40°16'N, long 38°46'E, alt. 1900 m, on *Fagus orientalis*, 11.09.2001, ANK 72.

Hypogymnia physodes (L.) Nyl.

Giresun: Şebinkarahisar; Kutluca Plateau, lat 40°11'N, long 38°48'E, alt. 2000 m, on *Pinus sylvestris*, 11.09.2001; Kirtepe-Yaygış, lat 40°13'N, long 38°40'E, alt. 2150 m, on *Abies nordmanniana* subsp. *nordmanniana*, 11.9.2001, ANK 17.

**Hypogymnia tubulosa* (Schaer.) Havaas

Giresun: Şebinkarahisar; Kutluca Plateau, lat 40°11'N, long 38°48'E, alt. 2000 m, on *Abies nordmanniana* subsp. *nordmanniana*, 11.09.2001, ANK 16.

****Hypogymnia vittata* (Ach.) Parr.**

Giresun: Şebinkarahisar; Doludere village, lat 40°16'N, long 38°39'E, alt. 1950 m., on *Pinus sylvestris*, 11.09.2001, ANK 73.

****Lecania fuscella* (Schaer.) Körb.**

Giresun: Şebinkarahisar; slope 4 km south of İncedere Plateau, lat 40°11'N, long 38°53'E, alt. 1750 m, on *Populus tremula*, 11.09.2001, ANK 74.

****Lecanora albella* (Pers.) Ach.**

Giresun: Şebinkarahisar; Arda Plateau, lat 40°18'N, long 38°46'E, alt. 1850 m, on *Cerasus mahaleb* var. *mahaleb*, 12.09.2001, ANK 75.

***Lecanora argentata* (Ach.) Malme**

Giresun: Şebinkarahisar; Kutluca Plateau, lat 40°11'N, long 38°48'E, alt. 2000 m, on *Abies nordmanniana* subsp. *nordmanniana*, 11.09.2001, ANK 76.

***Lecanora campestris* (Schaer.) Hue**

Giresun: Şebinkarahisar; Köroğlu Valley, lat 40°14'N, long 38°39'E, alt. 1750 m, on siliceous rocks, 12.09.2001, ANK 77.

****Lecanora carpinea* (L.) Vain.**

Giresun: Şebinkarahisar; Kutluca Plateau, lat 40°11'N, long 38°48'E, alt. 2000 m, on *Ulmus campestre*, 11.09.2001; Kabaktepe Hill, lat 40°13'N, long 38°38'E, alt. 1950 m, on *Carpinus betulus* sp., 11.09.2001, ANK 10.

***Lecanora chlarotera* Nyl.**

Giresun: Şebinkarahisar; slope 4 km south of İncedere Plateau, lat 40°11'N, long 38°53'E, alt. 1750 m, on *Abies nordmanniana* subsp. *nordmanniana*, 11.09.2001; Kabaktepe Hill, lat 40°13'N, long 38°38'E, alt. 1950 m, on *Abies nordmanniana* subsp. *nordmanniana*, 11.09.2001, ANK 78.

***Lecanora crenulata* s.l.**

Giresun: Şebinkarahisar; slope 4 km south of İncedere Plateau, lat 40°11'N, long 38°53'E, alt. 1750 m, on calcareous rocks, 11.09.2001, ANK 79.

****Lecanora polytropa* (Ehrh. ex Hoffm.) Rabenh.**

Giresun: Şebinkarahisar; Köroğlu Valley, lat 40°14'N, long 38°39'E, alt. 1750 m, on siliceous rocks, 12.09.2001, ANK 80.

****Lecanora saligna* (Schrad.) Zahlbr.**

Giresun: Şebinkarahisar; on top of and around Büyükçal Hill, lat 40°19'N, long 38°40'E, alt. 2000 m, on *Abies nordmanniana* subsp. *nordmanniana*, 12.09.2001, ANK 08.

****Lecanora varia* (Hoffm.) Ach.**

Giresun: Şebinkarahisar; on top of and around Büyükçal Hill, lat 40°19'N, long 38°40'E, alt. 2000 m, on *Abies nordmanniana* subsp. *nordmanniana*, 12.09.2001, ANK 09.

***Lecidella elaeochroma* (Ach.) M.Choisy**

Giresun: Şebinkarahisar; on top of and around Büyükçal Hill, lat 40°19'N, long 38°40'E, alt. 2000 m, on *Populus tremula*, 12.09.2001, ANK 09.

****Letharia vulpina* (L.) Hue**

Giresun: Şebinkarahisar; on top of and around Büyükçal Hill, lat 40°19'N, long 38°40'E, alt. 2000 m, on *Pinus sylvestris*, 12.09.2001; Giresun: Şebinkarahisar; Kutluca Plateau, lat 40°11'N, long 38°48'E, alt. 2000 m, on *Pinus sylvestris*, 11.09.2001; Arda Plateau, lat 40°18'N, long 38°46'E, alt. 1850 m, on *Pinus sylvestris*, 12.09.2001, ANK 21.

****Lobaria pulmonaria* (L.) Hoffm.**

Giresun: Şebinkarahisar; Yıldızlarla Hill, lat 40°14'N, long 38°48'E, alt. 1750 m, on *Abies nordmanniana* subsp. *nordmanniana*, 11.09.2001; Giresun: Şebinkarahisar; Doludere village, lat 40°16'N, long 38°39'E, alt. 1950 m, on calcareous rocks, 11.09.2001, ANK 81.

****Melanelia glabratula* (Lamy) O.Blanco et al.**

Giresun: Şebinkarahisar; Menekşeli-Ovacık, lat 40°16'N, long 38°46'E, alt. 1900 m, on *Abies nordmanniana* subsp. *nordmanniana*, 11.09.2001, ANK 24.

***Melanelia subaurifera* (Nyl.) O.Blanco et al.**

Giresun: Şebinkarahisar; Menekşeli-Ovacık, lat 40°16'N, long 38°46'E, alt. 1900 m, on *Abies nordmanniana* subsp. *nordmanniana*, 11.09.2001, ANK 26.

****Melanohalea elegantula* (Zahlbr.) O.Blanco et al.**

Giresun: Şebinkarahisar; Kirtepe-Yaygış, lat 40°13'N, long 38°40'E, alt. 2150 m, on *Abies nordmanniana* subsp. *nordmanniana*, 11.09.2001, ANK 22.

**Melanohalea exasperatula* (Nyl.) O.Blanco et al.

Giresun: Şebinkarahisar; Kirtepe-Yaygış, lat 40°13'N, long 38°40'E, alt. 2150 m, on *Abies nordmanniana* subsp. *nordmanniana*, 11.9.2001; Doludere village, lat 40°16'N, long 38°39'E, alt. 1950 m, on *Fagus orientalis*, 11.09.2001; slope 4 km south of İncedere Plateau, lat 40°11'N, long 38°53'E, alt. 1750 m, on mosses, 11.09.2001, ANK 23.

**Melanohalea infumata* (Nyl.) O.Blanco et al.

Giresun: Şebinkarahisar; on top of and around Büyüçal Hill, lat 40°19'N, long 38°40'E, alt. 2000 m, on *Abies nordmanniana* subsp. *nordmanniana*, 12.09.2001, ANK 25.

**Ochrolechia parella* (L.) A.Massal.

Giresun: Şebinkarahisar; slope 4 km south of İncedere Plateau, lat 40°11'N, long 38°53'E, alt. 1750 m, on *Carpinus betulus*, 11.09.2001, ANK 82.

Parmelia saxatilis (L.) Ach.

Giresun: Şebinkarahisar; Menekşeli-Ovacık, lat 40°16'N, long 38°46'E, alt. 1900 m, on mosses, 11.09.2001; Körülü valley, lat 40°14'N, long 38°39'E, alt. 1750 m, on siliceous rocks, 12.09.2001, ANK 27.

Parmelia sulcata Taylor

Giresun: Şebinkarahisar; Arda Plateau, lat 40°18'N, long 38°46'E, alt. 1850 m, on *Quercus infectoria* subsp. *infectoria*, 12.09.2001; Körülü valley, lat 40°14'N, long 38°39'E, alt. 1750 m, on *Quercus infectoria* subsp. *infectoria*, 12.09.2001, ANK 83.

Parmeliopsis ambigua (Wulfen) Nyl.

Giresun: Şebinkarahisar; Kutluca Plateau, lat 40°11'N, long 38°48'E, alt. 2000 m, on *Abies nordmanniana* subsp. *nordmanniana*, 11.09.2001; on top of and around Büyüçal Hill, lat 40°19'N, long 38°40'E, alt. 2000 m, on *Abies nordmanniana* subsp. *nordmanniana*, 12.09.2001, ANK 28.

Peltigera didactyla (With.) J.R.Laundon

Giresun: Şebinkarahisar; Körülü valley, lat 40°14'N, long 38°39'E, alt. 1750 m, on soil, 12.09.2001, ANK 42.

**Peltigera praetextata* (Flörke ex Sommerf.) Zopf

Giresun: Şebinkarahisar; Kabaktepe Hill, lat 40°13'N, long 38°38'E, alt. 1950 m, on soil, 11.09.2001, ANK 43.

Peltigera rufescens (Weiss) Humb.

Giresun: Şebinkarahisar; Menekşeli-Ovacık, lat 40°16'N, long 38°46'E, alt. 1900 m, on soil, 11.09.2001, ANK 84.

**Pertusaria amara* (Ach.) Nyl.

Giresun: Şebinkarahisar; on top of and around Büyüçal Hill, lat 40°19'N, long 38°40'E, alt. 2000 m, on *Quercus infectoria* subsp. *infectoria*, 12.09.2001, ANK 85.

Pertusaria lactea (L.) Arnold

Giresun: Şebinkarahisar; around Çakılıkaya, lat 40°12'N, long 38°50'E, alt. 2050 m, on siliceous rocks, 17.07.2002; Körülü valley, lat 40°14'N, long 38°39'E, alt. 1750 m, on siliceous rocks, 12.09.2001, ANK 86.

Phaeophyscia orbicularis (Neck.) Moberg

Giresun: Şebinkarahisar; on top of and around Büyüçal Hill, lat 40°19'N, long 38°40'E, alt. 2000 m, on soil, 12.09.2001, ANK 38.

Physcia stellaris (L.) Nyl.

Giresun: Şebinkarahisar; Doludere village, lat 40°16'N, long 38°39'E, alt. 1950 m., on *Abies nordmanniana* subsp. *nordmanniana*, 11.09.2001; Kabaktepe Hill, lat 40°13'N, long 38°38'E, alt. 1950 m, on *Quercus infectoria* subsp. *infectoria*, 11.09.2001, ANK 40.

**Physcia tenella* (Scop.) DC.

Giresun: Şebinkarahisar; on top of and around Büyüçal Hill, lat 40°19'N, long 38°40'E, alt. 2000 m, on *Abies nordmanniana* subsp. *nordmanniana*, 12.09.2001, ANK 39.

**Physconia distorta* (With.) J.R.Laundon

Giresun: Şebinkarahisar; Kutluca Plateau, lat 40°11'N, long 38°48'E, alt. 2000 m, on *Quercus infectoria* subsp. *infectoria*, 11.09.2001; Kabaktepe Hill, lat 40°13'N, long 38°38'E, alt. 1950 m, on *Quercus infectoria* subsp. *infectoria*, 11.09.2001; Arda plateau, lat 40°18'N, long 38°46'E, alt. 1850 m, on *Quercus infectoria* subsp. *infectoria*, 12.09.2001, ANK 87.

Plastismatia glauca (L.) W.L.Culb. & C.F.Culb.

Giresun: Şebinkarahisar; Kabaktepe Hill, lat 40°13'N, long 38°38'E, alt. 1950 m, on *Quercus infectoria* subsp. *infectoria*, 11.09.2001; Arda Plateau, lat 40°18'N, long 38°46'E, alt. 1850 m, on *Quercus infectoria* subsp. *infectoria*, 12.09.2001, ANK 88.

**Pleurosticta acetabulum* (Neck.) Elix & Lumbsch

Giresun: Şebinkarahisar; Arda Plateau, lat 40°18'N, long 38°46'E, alt. 1850 m, on *Fagus orientalis*, 12.09.2001, ANK 89.

Protoparmeliopsis muralis (Schreb.) M.Choisy

Giresun: Şebinkarahisar; on top of and around Büyüçal Hill, lat 40°19'N, long 38°40'E, alt. 2000 m, on siliceous rocks, 12.09.2001, ANK 11.

Pseudevernia furfuracea (L.) Zopf var. *ceratea* (Ach.) D.Hawksw.

Giresun: Şebinkarahisar; Kutluca Plateau, lat 40°11'N, long 38°48'E, alt. 2000 m, on *Pinus sylvestris*, 11.09.2001; Kirtepe-Yaygış, lat 40°13'N, long 38°40'E, alt. 2150 m, on *Pinus sylvestris*, 11.09.2001; Kabaktepe Hill, lat 40°13'N, long 38°38'E, alt. 1950 m, on *Pinus sylvestris*, 11.09.2001; Arda Plateau, lat 40°18'N, long 38°46'E, alt. 1850 m, on *Quercus infectoria* subsp. *infectoria*, 12.09.2001; Yıldızlarla Hill, lat 40°14'N, long 38°48'E, alt. 1750 m, on *Pinus sylvestris*, 11.09.2001, ANK 33.

Pseudevernia furfuracea (L.) Zopf var. *furfuracea*

Giresun: Şebinkarahisar; Kutluca Plateau, lat 40°11'N, long 38°48'E, alt. 2000 m, on *Pinus sylvestris*, 11.09.2001; Kabaktepe Hill, lat 40°13'N, long 38°38'E, alt. 1950 m, on *Pinus sylvestris*, 11.09.2001; Yıldızlarla Hill, lat 40°14'N, long 38°48'E, alt. 1750 m, on *Pinus sylvestris*, 11.09.2001, ANK 30.

Ramalina farinacea (L.) Ach.

Giresun: Şebinkarahisar; Kirtepe-Yaygış, lat 40°13'N, long 38°40'E, alt. 2150 m, on calcereous rocks, 11.09.2001, ANK 90.

**Ramalina fastigiata* (Pers.) Ach.

Giresun: Şebinkarahisar; Yıldızlarla Hill, lat 40°14'N, long 38°48'E, alt. 1750 m, on *Quercus infectoria* subsp. *infectoria*, 11.09.2001, ANK 91.

**Ramalina fraxinea* (L.) Ach.

Giresun: Şebinkarahisar; Köroğlu Valley, lat 40°14'N, long 38°39'E, alt. 1750 m, on siliceous rocks, 12.09.2001, ANK 92.

Ramalina pollinaria (Westr.) Ach.

Giresun: Şebinkarahisar; Köroğlu Valley, lat 40°14'N, long 38°39'E, alt. 1750 m, on siliceous rocks, 12.09.2001; Yıldızlarla Hill, lat 40°14'N, long 38°48'E,

alt. 1750 m, on *Abies nordmanniana* subsp. *nordmanniana*, 11.09.2001, ANK 93.

Rhizocarpon geographicum (L.) DC.

Giresun: Şebinkarahisar; around Çakılkaya, lat 40°12'N, long 38°50'E, alt. 2050 m, on calcareous rocks, 17.07.2002, ANK 94.

**Rinodina conradii* Körb.

Giresun; Şebinkarahisar; on top of and around Büyüçal Hill, lat 40°19'N, long 38°40'E, alt. 2000 m, on *Abies nordmanniana* subsp. *nordmanniana*, 12.09.2001, ANK 41.

**Rinodina pyrina* (Ach.) Arnold

Giresun: Şebinkarahisar; Kutluca Plateau, lat 40°11'N, long 38°48'E, alt. 2000 m, on *Abies nordmanniana* subsp. *nordmanniana*, 11.09.2001, ANK 95.

Tephromela atra (Huds.) Hafellner

Giresun: Şebinkarahisar; around Çakılkaya, lat 40°12'N, long 38°50'E, alt. 2050 m, on calcareous rocks, 17.07.2002 ANK 96.

**Umbilicaria cylindrica* (L.) Delise ex Duby

Giresun: Şebinkarahisar; around Çakılkaya, lat 40°12'N, long 38°50'E, alt. 2050 m, on siliceous rocks, 17.07.2002, ANK 97.

**Usnea hirta* (L.) Weber ex F.H.Wigg.

Giresun: Şebinkarahisar; Kabaktepe Hill, lat 40°13'N, long 38°38'E, alt. 1950 m, on *Pinus sylvestris*, 11.09.2001, ANK 98.

**Vulpicida pinastri* (Scop.) Mattson & Lai

Giresun: Şebinkarahisar; Kirtepe-Yaygış, lat 40°13'N, long 38°40'E, alt. 2000 m, on *Pinus sylvestris*, 12.9.2001, ANK 12.

Xanthomendoza ulophylloides (Räsänen) Söchting, Kärnefelt & S.Kondr.

Giresun: Şebinkarahisar; Köroğlu Valley, lat 40°14'N, long 38°39'E, alt. 1750 m, on calcareous rocks, 12.09.2001, ANK 99.

Xanthoria elegans (Link) Th.Fr.

Giresun: Şebinkarahisar; Köroğlu Valley, lat 40°14'N, long 38°39'E, alt. 1750 m, on calcareous rocks, 12.09.2001, ANK 100.

Discussion

There are also degraded forests in the study area; the dominant species of this vegetation type are *Quercus petrea* subsp. *iberica*, *Q. pubescens*, and *Q. infectoria* subsp. *infectoria*. All the vegetation types are under anthropogenic influence; therefore, lichen biodiversity in the study area is low.

Sarıçek Mountain is predominated by calcareous rocks. *Aspicilia calcarea*, *A. cinerea*, *Caloplaca biatorina*, *C. flavescens*, *Dermatocarpon intestiniforme*, and *Lecanora crenulata* are very abundant on calcareous rock in the study area. Siliceous rocks cover only a small part of Sarıçek Mountain and *Anaptychia ciliaris*, *Candelariella vitellina*, *Dimelaena oreina*, *Lecanora campestris*, *L. polytropa*, and *Parmelia saxatilis* are the dominant species in these areas. These findings are in agreement with the data in the literature (Purvis et al., 1992; Wirth, 1995).

The richest family according to the number of species is Parmeliaceae with 22 taxa, which comprises 27% of all collected species. For the other families, the number of taxa they have and their portion of all collected species are as follows: Lecanoraceae, 12 taxa (15%); Physciaceae, 8 taxa (10%); Teloschistaceae, 8 taxa (10%); Cladoniaceae, 4 taxa (5%); and Ramalinaceae, 4 taxa (5%).

Growth forms of the taxa differ. Of the 81 collected taxa, 68 are epiphytic, which is the dominant growth form. Because the dominant vegetation types in the study area are forests and degraded forests with high moss cover, which provide suitable habitats for lichens, *Abies*

sp. and *Pinus* sp. are preferred as a substrate by lichens. The other growth forms and their percent of total taxa are as follows: saxicolous, 31 taxa (28%); and terricolous, 12 taxa (11%).

There are *Pinus* sp., *Quercus* sp., and *Carpinus* sp. populations between 1750 and 1850 m, and so epiphytic lichens are abundant at these altitudes. Above 2000 m mountain steppic vegetation is observed and this area is dominated mostly by *Aspicilia* species.

The collection sites with the highest number of collected taxa are as follows: 1 (16 taxa), 7 (16 taxa), and 4 (15 taxa). Collection site 1 is situated at 2145 m and at site 7 rocky areas are dominant; the altitude and rocks prevent anthropogenic effects. Although there is a small village situated at collection site 4, it does not have a strong influence on the natural vegetation at this site, which also has a high number of taxa.

Kinalioğlu (2005) has identified 152 lichen species from northern Giresun Province. Our findings are similar, though for southern Giresun, and 48 taxa are new records for the province. The species found in the research area are common lichen species.

Air pollution and dry climate may be responsible for the rather low numbers of lichen taxa. The fact is that the effects of settlements vary at different parts of the mountain.

Acknowledgements

The authors thank Prof. Dr. Ayşen Özdemir Türk for her support in every aspect of the study.

References

- Akman Y (1990). *İklim ve Biyoiklim*. Ankara: Palme Yayınları Mühendislik Serisi, Turkey.
- Brummitt RK & Powell CE (1992). *Authors of Plant Names*. Kew: Royal Botanical Gardens Kew.
- Çobanoğlu G & Akdemir B (2004). Contribution to the lichen diversity of nature parks in Bolu and Çorum, Anatolia, Turkey. *Herzogia* 17: 129-136.
- Davis PH (1965). *Flora of Turkey and the East Aegean Islands*, Vol. 1. Edinburgh: Edinburgh University Press.
- Güvenç Ş (2002). Floristic records of lichens in Adana, Konya and Niğde Provinces. *Turk J Bot* 26: 175-180.
- Güvenç Ş & Öztürk Ş (2004). Lichen records from the alpine region of Uludağ (Olympus) Mountain in Bursa-Turkey. *Turk J Bot* 28: 299-306.
- John V & Breuss O (2004). Flechten der östlichen Schwarzmeer-Region in der Türkei (BLAM-Exkursion 1997). *Herzogia* 17: 137-155.
- Kinalioğlu K (2005). Lichens of Giresun District Giresun Province, Turkey. *Turk J Bot* 29: 417-423.
- Küçük M (1990). Giresun Adası'nın Floristik Yapısı. *Ormancılık Araştırma Enstitüsü Yayınları* 36: 2-58.
- Öztürk Ş, Güvenç Ş & Aydin S (2005). Floristic lichen records from Isparta and Burdur Provinces. *Turk J Bot* 29: 243-250.

- Özgen U, Aslan A & Terzi Z (2002). Phytochemical screening of some lichen species collected from Giresun province. *1st Int. Congr. on Chemistry of Natural Products*, 16-19 Oct. 2002-135, Trabzon.
- Purvis OW, Coppins BJ, Hawksworth DL, James PW & Moore DM (1992). *The Lichen Flora of Great Britain and Ireland*. London: Natural History Museum Publications in association with the British Lichen Society.
- Steiner J (1909). Ergebnisse einer bot. Reise in d. Pontische Randgebirge im Sandschak Trapezunt. *Lichenes*. In: D.H.F. Handel-Mazzetti, *Annal Naturh Hofmus Wien* 23: 107-123.
- Süleyman H, Yıldırım D, Aslan A, Göçer F, Gepdiremen A & Güvenalp Z (2002). An investigation of the anti-inflammatory effects of an extract from *Cladonia rangiformis* Hoffm. *Biol Pharm Bull* 25: 10-13.
- Yazıcı K & Aslan A (2005). Six new lichen records from Turkey. *Mycotaxon* 93: 359-363.
- Wirth V (1995). *Die Flechten Baden – Württembergs*. Teil 1-2. Stuttgart: Ulmer.
- Yıldız A & John V (2002). Additional lichen records from Kastamonu Province (Turkey). *Fl Medit* 12: 315-322.
- Yıldız A, John V & Yurdakulol E (2002). Lichens from the Çangal Mountains (Sinop, Turkey). *Cryptogamie Mycol* 23: 81-88.